Brain Research, 409 (1987) 409–411
Elsevier

Author Index

Acker, H. and Eyzaguirre, C., Light absorbance changes in the mouse carotid body during hypoxia and cyanide poisoning, 380
Adrien, J., see Davenne, D., 1
Aikawa, E., see Kawano, J., 391
Alavi, A., see Kushner, M.J., 79
Alford, S. and Williams, T.L., Inhibitory synaptic input to edge cells during fictive locomotion, 139
Atsumi, S. and Sakamoto, H., Enkephalin-like immunoreactive axon terminals make synapses with α-motoneurons in the chicken, 187

Berglund, S., see Brismar, T., 227
Bernstein, J.J., see Connor, J.R., 62
Bignami, A. and Clark, K., Non-phosphorylated and phosphorylated neurofilaments in hypothyroid rat cerebellum, 143
Bitar, M.S., Koulu, M. and Linnoila, M., Diabetes-induced changes in monoamine concentrations of rat hypothalamic nuclei, 236
Brennan, T.J., see Girardot, M.-N., 19
Bressler, S.L., Relation of olfactory bulb and cortex. I. Spatial variation of bulbocortical interdependence, 285
Bressler, S.L., Relation of olfactory bulb and cortex. II. Model for driving of cortex by bulb, 294
Brismar, T., Hildebrand, C. and Berglund, S., Voltage-clamp analysis of nodes of Ranvier in regenerated rat sciatic nerve, 227
Brown, M.W., Wilson, F.A.W. and Riches, I.P., Neuronal evidence that inferomedial temporal cortex is more important than hippocampus in certain processes underlying recognition memory, 158
Bryld, E., Zeeberg, I., Gjerris, A., Werdelin, L. and Rehfeld, J.F., Increased cerebrospinal fluid concentrations of C- but not N-terminal cholecystokinin fragments in multiple sclerosis, 364
Burton, P.R., Microtubules of frog olfactory axons: their length and number/axon, 71

Carvey, P.M., Kao, L.C. and Klawans, H.L., The effect of bilateral kainic acid-induced lateral habenula lesions on dopamine-mediated behaviors, 193
Caspers, M.L., Schwartz, R.D., Labarca, R. and Paul, S.M., Autoradiographic visualization and characterization of [^{3}H]ouabain binding to the Na^+,K^+-ATPase of rat brain and pineal, 335
Clark, K., see Bignami, A., 143
Cobbett, P. and Mason, W.T., Whole cell voltage clamp recordings from cultured neurons of the supraoptic area of neonatal rat hypothalamus, 175
Coggeshall, R.E., see Jenq, C.-B., 250
Cohen, G., see Slivka, A., 275
Cohen-Becker, I., see Osterburg, H.H., 31
Coleman, P.D., see Flood, D.G., 88
Collier, T.J., Quirk, G.J. and Routtenberg, A., Separable roles of hippocampal granule cells in forgetting and pyramidal cells in remembering spatial information, 316
Connor, J.R. and Bernstein, J.J., Astrocytes in rat fetal cerebral cortical homografts following implantation into adult rat spinal cord, 62

D'Angio, M., Serrano, A., Rivy, J.P. and Scatton, B., Tail-pinch stress increases extracellular DOPAC levels (as measured by in vivo voltammetry) in the rat nucleus accumbens but not frontal cortex: antagonism by diazepam and zolpidem, 169
Dann, R., see Kushner, M.J., 79
Dark, J., Dark, K.A. and Zucker, I., Long day lengths increase brain weight and DNA content in the meadow vole, *Microtus pennsylvanicus*, 302
Dark, K.A., see Dark, J., 302
Davenne, D. and Adrien, J., Lesion of the ponto-geniculo-occipital pathways in kittens. I. Effects on sleep and on unitary discharge of the lateral geniculate nucleus, 1

Emson, P.C., see Kawai, Y., 371
Evans, D. and Smith, J.C., Seizure activity and cortical spreading depression monitored by an extrinsic potential-sensitive molecular probe, 350
Eyzaguirre, C., see Acker, H., 380

Fifková, E., see Scheetz, A.J., 329
Finch, C.E., see Osterburg, H.H., 31
Flood, D.G., Guarnaccia, M. and Coleman, P.D., Dendritic extent in human CA_{2-3} hippocampal pyramidal neurons in normal aging and senile dementia, 88
Foreman, R.D., see Girardot, M.-N., 19

Gabriel, M. and Sparenborg, S., Posterior cingulate cortical lesions eliminate learning-related unit activity in the anterior cingulate cortex, 151
Gavish, M., Weizman, A., Youdim, M.B.H. and Okun, F., Regulation of central and peripheral benzodiazepine receptors in progesterone-treated rats, 386
Gehlert, D.R. and Wamsley, J.K., Quantitative autoradiography of α_2 agonist binding sites in the spontaneously hypertensive rat brain, 308
Getting, P.A., see Richerson, G.B., 128
Girardot, M.-N., Brennan, T.J., Martindale, M.E. and Foreman, R.D., Effects of stimulating the subcoeruleus–parabrachial region on the non-noxious and noxious responses of T_1–T_5 spinothalamic tract neurons in the primate, 19
Girgis, S., see Kawai, Y., 371
Gjerris, A., see Bryld, E., 364
Groves, P.M., see Klemfuss, H., 197
Guarnaccia, M., see Flood, D.G., 88

Hama, K., see Kosaka, T., 403
Hamaoka, Y., see Kosaka, T., 403
Harrison, B.M., Schwann cells divide in a demyelinating lesion of the central nervous system, 163
Hasegawa, A., Ohtsubo, K. and Mori, W., Pineal gland in old age; quantitative and qualitative morphological study of 168 human autopsy cases, 343
Heizmann, C.W., see Kosaka, T., 403

Henke, P.G., see Ray, A., 398
Hildebrand, C., see Brismar, T., 227
Hillyard, C.J., see Kawai, Y., 371
Hoffmann, K.-P., see Horn, A.K.E., 133
Hoogendijk, J.E., see Roozendaal, B., 259
Horn, A.K.E. and Hoffmann, K.-P., Combined GABA-immunocytochemistry and TMB–HRP histochemistry of pretectal nuclei projecting to the inferior olive in rats, cats and monkeys, 133

Itaya, S.K., Anterograde transsynaptic transport of WGA-HRP in rat olfactory pathways, 205
Ito, K. and McCarley, R.W., Physiological studies of brainstem reticular connectivity. I. Responses of mPRF neurons to stimulation of bulbar reticular formation, 97
Ito, K., see McCarley, R.W., 111

Jenq, C.-B. and Coggeshall, R.E., Single vs multiple somatic nerve crushes in the rat, 250

Kannan, H., Kasai, M., Osaka, T. and Yamashita, H., Neurons in the paraventricular nucleus projecting to the median eminence: a study of their afferent connections from peripheral baroreceptors, and from the A_1-catecholaminergic area in the ventrolateral medulla, 358
Kao, L.C., see Carvey, P.M., 193
Kasai, M., see Kannan, H., 358
Kastin, A.J., see Zadina, J.E., 10
Kawai, Y., Emson, P.C., Hillyard, C.J., Girgis, S., MacIntyre, I., Oertel, W.H. and Tohyama, M., Immunohistochemical evidence for the coexistence of calcitonin gene-related peptide and glutamate decarboxylase-like immunoreactivities in the Purkinje cells of the rat cerebellum, 371
Kawano, J. and Aikawa, E., Regional distribution of arylsulfatase C and estrone-sulfate sulfatase activities in rat brain and hypophysis, 391
Kigoshi, S., see Yamanaka, K., 395
Klawans, H.L., see Carvey, P.M., 193
Klemfuss, H., Young, S.J. and Groves, P.M., Do antidromic latency jumps indicate axonal branching in nigrostriatal and hypothalamo-neurohypophysial neurons?, 197
Kosaka, T., Heizmann, C.W., Tateishi, K., Hamaoka, Y. and Hama, K., An aspect of the organizational principle of the γ-aminobutyric acidergic system in the cerebral cortex, 403
Koulu, M., see Bitar, M.S., 236
Kushner, M.J., Schwartz, R., Alavi, A., Dann, R., Rosen, M., Silver, F. and Reivich, M., Cerebral glucose consumption following verbal auditory stimulation, 79

Labarca, R., see Caspers, M.L., 335
LeRoith, D., see Waldbillig, R.J., 215
Li, Y.-W., see Su, H.-S., 367
Linnoila, M., see Bitar, M.S., 236

MacIntyre, I., see Kawai, Y., 371
Manasco, P.K., see Zadina, J.E., 10
Markham, J.A., see Scheetz, A.J., 329
Martindale, M.E., see Girardot, M.-N., 19
Mason, W.T., see Cobbett, P., 175
McCarley, R.W., Ito, K. and Rodrigo-Angulo, M.L., Physiological studies of brainstem reticular connectivity. II. Responses of mPRF neurons to stimulation of mesencephalic and contralateral pontine reticular formation, 111
McCarley, R.W., see Ito, K., 97
Mirmiran, M., see Roozendaal, B., 259
Morgan, D.G., see Osterburg, H.H., 31
Mori, W., see Hasegawa, A., 343
Muramatsu, I., see Yamanaka, K., 395
Mytilineou, C., see Slivka, A., 275

Nastiuk, K.L., see Zadina, J.E., 10
Nilsson, G.E. and Tottmar, O., Effects of biogenic aldehydes and aldehyde dehydrogenase inhibitors on rat brain tryptophan hydroxylase activity in vitro, 374
Nilsson, G.E., Tottmar, O. and Wahlström, G., Effects of aldehyde dehydrogenase inhibitors on hexobarbital sensitivity and neuroamine metabolism in rat brain, 265
Oertel, W.H., see Kawai, Y., 371
Ohtsubo, K., see Hasegawa, A., 343
Okun, F., see Gavish, M., 386
Osaka, T., see Kannan, H., 358
Osterburg, H.H., Telford, N.A., Morgan, D.G., Cohen-Becker, I., Wise, P.M. and Finch, C.E., Hypothalamic monoamines and their catabolites in relation to the estradiol-induced luteinizing hormone surge, 31
Oyama, Y., Actions of convulsants, 4-aminopyridine and pentylenetetrazole, on the transient outward current of single isolated nodose ganglion neurons, 243

Paul, S.M., see Caspers, M.L., 335
Peng, Z.-C., see Su, H.-S., 367
Pignatiello, M.F., see Zadina, J.E., 10
Pritz, M.B. and Stritzel, M.E., Percentage of intrinsic and relay cells in a thalamic nucleus projecting to general cortex in reptiles, *Caiman crocodilus*, 146

Quirk, G.J., see Collier, T.J., 316

Ray, A., Henke, P.G. and Sullivan, R.M., The central amygdala and immobilization stress-induced gastric pathology in rats: neurotensin and dopamine, 398
Rehfeld, J.F., see Bryld, E., 364
Reivich, M., see Kushner, M.J., 79
Richerson, G.B. and Getting, P.A., Maintenance of complex neural function during perfusion of the mammalian brain, 128
Riches, I.P., see Brown, M.W., 158
Rivy, J.P., see D'Angio, M., 169
Rodrigo-Angulo, M.L., see McCarley, R.W., 111
Roozendaal, B., Van Gool, W.A., Swaab, D.F., Hoogendijk, J.E. and Mirmiran, M., Changes in vasopressin cells of the rat suprachiasmatic nucleus with aging, 259
Rosen, M., see Kushner, M.J., 79
Routtenberg, A., see Collier, T.J., 316

Sakamoto, H., see Atsumi, S., 187
Scatton, B., see D'Angio, M., 169
Scheetz, A.J., Markham, J.A. and Fifková, E., The effect of chronic ethanol consumption on the fine structure of the CA_1 stratum oriens in short-sleep and long-sleep mice: short-term and long-term exposure, 329
Schwartz, R., see Kushner, M.J., 79
Schwartz, R.D., see Caspers, M.L., 335
Serrano, A., see D'Angio, M., 169
Shinoda, K., Tohyama, M. and Shiotani, Y., Hippocampofugal γ-aminobutyric acid (GABA)-containing neuron system in the rat: a study using a double-labeling method that combines retrograde tracing and immunocytochemistry, 181

Shiotani, Y., see Shinoda, K., 181
Silver, F., see Kushner, M.J., 79
Slivka, A., Mytilineou, C. and Cohen, G., Histochemical evaluation of glutathione in brain, 275
Smith, J.C., see Evans, D., 350
Sparenborg, S., see Gabriel, M., 151
Strausbaugh, L.J., Intracarotid infusions of protamine sulfate disrupt the blood–brain barrier of rabbits, 221
Stritzel, M.E., see Pritz, M.B., 146
Su, H.-S., Peng, Z.-C. and Li, Y.-W., Distribution of catecholamine-containing cell bodies in the human diencephalon, 367
Sullivan, R.M., see Ray, A., 398
Swaab, D.F., see Roozendaal, B., 259

Taira, K., Characteristics of periodontal mechanosensitive neurons in the first somatosensory cortex of the cat, 52
Taira, K., The representation of the oral structures in the first somatosensory cortex of the cat, 41
Tateishi, K., see Kosaka, T., 403
Telford, N.A., see Osterburg, H.H., 31
Tohyama, M., see Kawai, Y., 371
Tohyama, M., see Shinoda, K., 181
Tottmar, O., see Nilsson, G.E., 265
Tottmar, O., see Nilsson, G.E., 374

Van Gool, W.A., see Roozendaal, B., 259

Wahlström, G., see Nilsson, G.E., 265
Waldbillig, R.J. and LeRoith, D., Insulin receptors in the peripheral nervous system: a structural and functional analysis, 215
Wamsley, J.K., see Gehlert, D.R., 308
Weizman, A., see Gavish, M., 386
Werdelin, L., see Bryld, E., 364
Williams, T.L., see Alford, S., 139
Wilson, F.A.W., see Brown, M.W., 158
Wise, P.M., see Osterburg, H.H., 31

Yamanaka, K., Muramatsu, I. and Kigoshi, S., Muscarinic agonist binding in rat brain following chronic nicotine treatment, 395
Yamashita, H., see Kannan, H., 358
Youdim, M.B.H., see Gavish, M., 386
Young, S.J., see Klemfuss, H., 197

Zadina, J.E., Kastin, A.J., Manasco, P.K., Pignatiello, M.F. and Nastiuk, K.L., Long-term hyperalgesia induced by neonatal β-endorphin and morphiceptin is blocked by neonatal Tyr-MIF-1, 10
Zeeberg, I., see Bryld, E., 364
Zucker, I., see Dark, J., 302